

Flora of Nepal

नेपालका वनस्पति

Ehretiaceae

Ehretiaceae: Webedition 1 (2023)

<https://data.rbge.org.uk/publications/FloraofNepal/library/Ehretiaceae/1>

Editors

Mark F. Watson, Shinobu Akiyama, Hiroshi Ikeda, Colin A. Pendry, Keshab R. Rajbhandari, Krishna K. Shrestha

Authored by

Pendry C.A.

Genera in this account

Ehretia (p.1)

Published on 02 June 2023 by Royal Botanic Garden Edinburgh, 20a Inverleith Row, Edinburgh, EH2 5LR, UK

This PDF Webedition will be permanently available and citable using the URL specified at the top of this page. Family accounts have numbered pages so individual pages can be cited. Previous and future editions can also be accessed via the project website (www.floraofnepal.org/PDFLibrary).

© text and images, except where individually credited,

Nepal Academy of Science and Technology, Khumaltar, Lalitpur, Kathmandu, Nepal

Royal Botanic Garden Edinburgh, 20a Inverleith Row, Edinburgh, EH3 5LR, UK

The Society of Himalayan Botany, University Museum, University of Tokyo, Hongo 7-3-1, Tokyo 113-0033, Japan

All rights reserved. 2017

Every effort has been made to trace holders of copyright in text and illustrations. Should there be any inadvertent omissions or errors the publishers will be pleased to correct them for future editions.

Appendices

1: Illustration plates

2: Format, abbreviations and categories used in Flora of Nepal

See printed volumes of the Flora of Nepal (www.rbge.org.uk/publications/floraofnepal) and the project website (www.floraofnepal.org) for further information on the Flora of Nepal and acknowledgement of the institutes and people involved with this international collaborative project.



Royal
Botanic Garden
Edinburgh



Ehretiaceae C.Mart.

C.A. Pendry

Trees. Leaves petiolate, alternate, margins serrate or entire, venation pinnate. Inflorescences terminal or axillary, dichotomous cymes, sometimes with spicate terminal branches, bracteate or ebracteate. Flowers bisexual, regular. Calyx cup-shaped, with a short tube and 5 lobes, not accrescent. Corolla 5-merous, tube longer or shorter than the reflexed or spreading lobes. Stamens 5, inserted at the throat of the corolla tube, filaments glabrous, anthers medifixed, exserted. Ovary 4-locular, ovules 1 per locule. Style terminal, 2-cleft with short lobes. Fruit a 4-celled drupe with a hard endocarp breaking up into two 2-seeded or four 1-seeded pyrenes.

Worldwide nine genera and about 150 species in tropical and warm temperate regions. One genus and five species in Nepal.

1. *Ehretia* P. Browne, Civ. Nat. Hist. Jam.: 168 (1756).

Description as for Ehretiaceae.

Key to Species


- | | | |
|----|-------------------------------------------------------------------------------------------------------------------------------|--------------------------|
| 1a | Leaf margins serrate | 2 |
| b | Leaf margins entire | 3 |
| 2a | Flowers 4–5 mm across, corolla tube included within the calyx. Leaves elliptic to slightly obovate, 6–13 × 1.5–4 cm | 1. <i>E. acuminata</i> |
| b | Flowers 6–9 mm across, corolla tube exserted from the calyx. Leaves broadly ovate to elliptic, 8–17(–28) × 5–12(–18) cm | 2. <i>E. macrophylla</i> |
| 3a | Corolla subrotate, lobes spreading | 5. <i>E. aspera</i> |
| b | Corolla funnel-shaped, lobes reflexed | 4 |
| 4a | Corolla glabrous outside, lobes 4–5 mm. Stigma lobes 0.8–1 mm | 3. <i>E. psilosiphon</i> |
| b | Corolla minutely puberulent outside, lobes to 3.5 mm. Stigma lobes 1.8–2.5 mm | 4. <i>E. wallichiana</i> |

1. *Ehretia acuminata* R.Br., Prodr. Fl. Nov. Holland.: 497 (1810).

Ehretia acuminata var. *serrata* (Roxb.) I.M.Johnst.; *E. pyrifolia* D. Don.; *E. serrata* Roxb.

Tree to 15 m. Twigs glabrous. Petioles 1–3 cm. Leaves elliptic to slightly obovate, 6–13 × 1.5–4 cm, base cuneate, apex acuminate, margins serrate, veins 5–6 pairs, glabrous above and below. Inflorescences terminal and axillary, paniculate, conical, 4–15 × 2–12 cm, axes sparsely pubescent. Inflorescence bracts caducous. Calyx cup-shaped, 2 mm, tube 1 mm, glabrous, lobes rounded, 1 mm, glabrous, minutely ciliate. Flowers 4–5 mm across, corolla white, subglabrous, tube 1–1.5 mm, lobes spreading 2 mm. Free part of stamens 2.5–3 mm, anthers 1–1.3 mm. Ovary green, 1 mm, style 2 mm, lobes 0.5 mm. Fruit orange, ca. 0.5 cm.

Distribution: Nepal, E Himalaya and Assam-Burma.



Altitudinal range: (400–)1,000–2,200 m.


Ecology: Open hillsides.

Flowering: April–June. **Fruiting:** (June)July–August.

2. *Ehretia macrophylla* Wall., Fl. Ind. (Roxburgh) 2: 343 (1824).

Tree to 10 m. Twigs glabrous or with scattered, sparse long hairs. Petioles 1.4–2.5 cm. Leaves broadly ovate to elliptic, 8–17 × 5–12 cm, base rounded to truncate or cuneate, apex acuminate, margins finely, sharply serrate, veins 6–7 pairs, scabrid above, glabrous below. Inflorescences terminal, rounded, 6–12 × 7–15 cm, with flowers in dense, cymose clusters on long branches, axes long pubescent. Inflorescence bracts to 1 cm, persistent. Calyx cup-shaped, 3–4 mm, tube 1–1.5 mm, glabrous, lobes narrowly triangular, 2.5–3 mm, shortly pubescent. Flowers 6–9 mm across, corolla white or cream, subglabrous, tube 5–6.5 mm, lobes reflexed, 3–4 mm. Free part of stamens 4–5 mm, anthers 1.4 mm. Ovary green, 1 mm, style 8 mm, lobes 1–1.5 mm. Fruit yellow, ca. 1 cm.

Distribution: Nepal, W Himalaya, E Himalaya, Tibetan Plateau, Assam-Burma, E Asia and SE Asia.



Altitudinal range: 1,500–2,500 m.

Ecology: *Pinus roxburghii* and broadleaved forest, open hillsides.


Flowering: April–May. **Fruiting:** (May–)June–August.

Flowers fragrant.

3. *Ehretia psilosiphon* R.R.Mill, Edinburgh J. Bot. 53(2): 260 (1996).

Tree to 7 m. Twigs glabrous. Petioles 1–2 cm. Leaves elliptic to slightly ovate or slightly obovate, 5–12 × 2–5.5 cm, base narrowly cuneate, apex acuminate, margins entire, veins 5–6 pairs, glabrous above and below. Inflorescences terminal, sometimes on short side branches, dichotomous below, spicate above, 5–12 × 5–12 cm, axes glabrous. Bracts absent. Calyx cup-shaped, glabrous, 2.5 mm, tube 1 mm, lobes triangular, 1.5 mm, minutely ciliate. Flowers 5–7 mm across, corolla white, funnel-shaped, glabrous, tube 6–7 mm, lobes reflexed, 4–5 mm. Free part of stamens 6 mm, anthers 2.2–2.5 mm. Ovary 1–1.5 mm, style 7–8 mm, lobes 0.8–1 mm. Fruit not seen.

Distribution: Nepal and E Himalaya.



Altitudinal range: 200–700 m.

Ecology: Mixed moist tropical forest.

Flowering: March–April. **Fruiting:** May.


Flowers said to have an unpleasant fragrance.

4. *Ehretia wallichiana* Hook.f. & Thomson ex. C.B.Clarke, Fl. Brit. India 4[10]: 143 (1883).

Cordia acuminata Wall.

Tree to 15 m. Twigs glabrous. Petioles 1–2(–3) cm. Leaves oblong-elliptic to slightly ovate, 5–15 × 3–7 cm, base cuneate, apex shortly acuminate, margins entire, veins 6–7 pairs, glabrous above and below. Inflorescences axillary in axils of previous year's fallen leaves, globose, repeatedly dichotomous, on a 1–2 cm peduncle, axes sparsely pubescent. Bracts absent. Calyx cup-shaped, glabrous, 2.5–3 mm, tube 1 mm, lobes triangular, 1.5–2 mm, minutely ciliate. Flowers 6–9 mm across, corolla white, funnel-shaped, minutely puberulent outside, glabrous within, tube 5.5–6 mm, lobes reflexed, 2.5–3.5 mm. Free part of stamens 4.5–6 mm, anthers 1.5 mm. Ovary 1–1.5 mm, style 7–9 mm, lobes 1.8–2.5 mm. Fruit red- or yellow-green, globose, ca. 1 cm.

Distribution: Nepal, E Himalaya and Assam-Burma.



Altitudinal range: 1,300–1,700 m.

Ecology: Mixed moist tropical and lower temperate forest.

Flowering: April–May. **Fruiting:** May–July.


Flowers sweetly fragrant.

5. *Ehretia aspera* Willd., Phytographia: 4 (1794).

Ehretia laevis Roxb.; *E. punctata* Roth ex. Roem. & Schult.

Tree to 15 m. Twigs glabrous. Petioles 1–1.5(–2) cm. Leaves elliptic-oblong to slightly ovate or slightly obovate, 4–14 × 3–7 cm, base cuneate, apex shortly acuminate, margins entire, veins 4–6 pairs, glabrous above and below. Inflorescences terminal and axillary, dichotomous below, with spicate branches above, 5–10 × 5–8 cm, axes puberulent to subglabrous. Bracts caducous. Calyx shallowly cup-shaped, puberulent to subglabrous, 1.5–2.5 mm, tube 0.5–0.7 mm, lobes triangular, 1–1.5 mm, minutely ciliate. Flowers 6–8 mm across, corolla white, subrotate, glabrous, tube 1.5–2 mm, lobes spreading, 2.5–3 mm. Free part of stamens 2–2.5 mm, anthers 1 mm. Ovary 1 mm, style 1.5–2 mm, lobes 0.5–1 mm. Fruit orange, ca. 0.5 cm.

Distribution: Nepal, W Himalaya, E Himalaya, Assam-Burma, S Asia and SE Asia.



Altitudinal range: 100–1,100 m.

Ecology: Riverine forest, sal forest, secondary forest, dry slopes.

Flowering: February–June. **Fruiting:** April–June.

How to use this pdf web edition

This Web-edition pdf document forms part of a set of Flora accounts for families and genera that have been finalized, including those in volumes yet to be printed. These pdf documents are made accessible via the *Flora of Nepal* website (www.floraofnepal.org) and will be periodically updated in numbered versions, permanently available and citable.

Flora of Nepal takes an innovative approach to Flora writing, with an underlying data base system managing the *Flora of Nepal Knowledge Base* from which the printed volumes and the 'online Flora' (www.floraofnepal.org) are generated. The Internet-accessible dataset augments the printed Flora by presenting all herbarium specimen data, detailed taxonomic information (such as full nomenclatural references and typification), distribution maps with point occurrences and images used when preparing the Flora. Much of this information is accumulated as a normal part of taxonomic working practices when undertaking a floristic revision, but it is usually lost to a wider audience as it is rarely included in the traditional printed Flora.

Flora of Nepal includes all native and fully naturalized vascular plants recorded within the political borders of Nepal, including brief references to agricultural and horticultural plants as appropriate. For pragmatic reasons the arrangement of families in the printed *Flora of Nepal* follows a modified Englerian sequence, closely following that of the *Flora of China* and, to a lesser extent, the *Flora of Bhutan*.^{1,2} In recent years the world view on the arrangement of families has radically changed following overwhelming phylogenetic evidence. The emergent family-level classification, now in its third iteration as APG IV, is reasonably stable and widely accepted.³ It has not been possible to alter the family sequence in *Flora of Nepal* printed volumes midway through the project, but as the data are stored separately in a database, the families can be reorganized electronically at a later date to reflect alternative classifications. Circumscription of families and genera, however, generally does follow a contemporary understanding of their relationships, except where group experts advise otherwise. Genera and species are treated in taxonomic order, or if there is disagreement then morphologically similar species are usually grouped together or occasionally listed alphabetically. Intraspecific taxa are always presented in alphabetical order. Intermediate ranks, such as subfamily, tribe, subgenus, section and series, are only used when they are useful in the treatment of large families or genera.

Information on nomenclature and classification is given for all accepted scientific names and synonyms pertaining to Nepal and nearby regions. Emphasis is given to those names listed in the primary checklists for Nepal: *Enumeration of the Flowering Plants of Nepal*,⁴ *Annotated Checklist of the Flowering Plants of Nepal*,⁵ and *Flowering Plants of Nepal (Phanerogams)*.⁶ At the generic level, synonyms widely used in the Asian literature are included. Full bibliographic citation with authorship is given for all accepted names and their basionyms at the rank of genus and below. As far as possible, the bibliographic citations of all accepted names and their basionyms have been verified with the original literature. The basionym precedes all other synonyms, which are listed alphabetically. Misapplied names (misidentifications encountered in the literature) are not included in synonymy, but are discussed in the supporting information at the end of a taxon. Authors of plant names follow the standard forms given in *Authors of Plant Names* and its continuously updated online supplement (www.ipni.org).⁷ Bibliographic references are given using the standard abbreviations in BPH-2 for serial publications (journals and periodicals) and in TL-2 (and its supplements) for books.^{8,9} In some cases books were published in several fascicles on different dates, sometimes over different years, but not indicated as such in the printed work. Date of publication is critical for establishing nomenclatural priority, and so it is important to be precise when citing names published in such works. The fascicle composition and publication dates of these often complex cases are clearly explained in TL-2, but the standard abbreviation does not differentiate between them. In these instances the TL-2 abbreviation has been amended with brackets to clearly indicate which fascicle is being referred to, for example Wallich, N., Pl. As. Rar. 2[8]. 1831. Books and periodicals not included in these two standard references have been abbreviated according to the recommendation in Appendix A of BPH-2.

Where a taxon has a widely recognized local name this is given in Devanagari script, followed by its transliteration into the Latin alphabet and the language of the vernacular name in parentheses '()'. One local name is given in the printed Flora, whereas multiple alternative vernacular names in different languages may be included in the *Flora of Nepal Knowledge Base* and made available online. Separate indexes to vernacular names in Devanagari, their Latin transliterations and scientific names are included at the end of each volume.

Descriptions are given for all taxa (family, genus, species, infraspecies and occasionally intermediate ranks) and wherever possible are based on primary observations and measurements made on specimens from Nepal. If no such material was available to authors, descriptions are taken from specimens from adjacent countries or secondary sources, and annotated as such. Most descriptions are about 150 words long, but exceptionally they are shorter or longer depending on the complexity of the taxon being described. For species with more than one infraspecific taxon, a full description is given for the species and short diagnoses for the lower taxa. Descriptions aim to be consistent and parallel between taxa of the same rank within a higher taxon. Authors were asked to standardize descriptive terms using the definitions given in *Plant Identification Terminology*.¹⁰ If a single measurement is given it refers to

length, and if width is also given it is in the format length × width. Ranges are separated by an en-dash (–) and discontinuous states by the word 'or'. Exceptional measurements are given in parentheses '()'. Taxon statistics and short statements on worldwide distribution are provided for families and genera, with summary statistics of lower taxa represented in Nepal.

Identification keys are dichotomous and presented in a bracketed format, with all elements strictly parallel between the two leads of each couplet. Keys are artificial and not intended to reflect any taxonomic classification. There is usually a single key to genera within a family, combining flowering, fruiting and vegetative characters, but where this is unwieldy separate keys are given for flowering and fruiting material (e.g. Cruciferae, Rosaceae). Keys are also given for species within a genus and taxa within a species. Figures are provided to aid identification by illustrating the diagnostic characters of each family and genus, and for large genera variation in major morphological features is represented.

The geographic distribution within Nepal is indicated for each species and infraspecific taxon at the political district level by a shaded distribution map. The distribution maps are evidence-based, produced from the *Flora of Nepal Knowledge Base* using locality information taken from authenticated herbarium specimens and records of plants *in situ* made by credible observers. Ideally all specimens identified by authors should be geo-referenced and databased when they are preparing *Flora of Nepal* accounts, but where this is not possible a minimum of one specimen per district is required. Sometimes the distribution of a species is greater than the sum of the distribution maps of its infraspecific taxa. This is a result of some herbarium specimens only being identifiable to species level. Occasionally species are known only from poorly localised collections, especially those from the early 19th century. For example, Wallich often only gave 'Nepalia' as the locality for many of his 1820–1821 collections. These specimens are most likely to have come from the Kathmandu Valley, known as the 'Nepal Valley' or just 'Nepal' at that time, but they might also have been collected during his inward and outward journeys from India via Hetauda, or by pilgrims going north to 'Gossainthan' (Gossainkund). It is therefore impossible to be sure of the correct district and in such cases this is noted in the supporting information and the map omitted. The *Flora of Nepal* website gives access to the underlying distribution and specimen information through an interactive dot map plotting all geo-referenced occurrence records and a listing of all material recorded.

Distribution for species and infraspecific taxa occurring outside Nepal is indicated by a list of geographical regions, with the resolution becoming coarser with increasing distance from Nepal. In order to utilise information contained within other published Floras these areas are defined according to political borders, with countries or provinces grouped to form regions that have some underlying biogeographic basis. For example, although the Tibetan Plateau extends into parts of Sichuan and Yunnan, we limit it to Xizang and Qinghai. *Flora of Nepal* takes no stance on any politically disputed border areas and is following the current international mapping convention of using the 'lines of control' to delineate its regions. The names used for the regions are intended to be descriptive and non-political. The regions are:

<i>W Himalaya</i>	India (Jammu & Kashmir, Himachal Pradesh, Uttarakhand), northern Pakistan (Khyber Pakhtunkhwa, previously known as North West Frontier Province).
<i>E Himalaya</i>	Sikkim, Darjeeling, Bhutan, NE India (Arunachal Pradesh).
<i>Tibetan Plateau</i>	China (Xizang, Qinghai).
<i>Assam-Burma</i>	Assam, Nagaland, Manipur, Myanmar
<i>S Asia</i>	Eastern Pakistan (Punjab, Sind, Islamabad), Peninsular India, Sri Lanka, Bangladesh, Maldives.
<i>E Asia</i>	China (excluding Xizang, Xinjiang, Qinghai), Korea, Japan, Taiwan.
<i>SE Asia</i>	Thailand, Laos, Cambodia, Vietnam, Malaysia, Indonesia, Philippines, New Guinea.
<i>N Asia</i>	China (Xinjiang), Russia, Mongolia.
<i>C Asia</i>	Kazakhstan, Uzbekistan, Turkmenistan, Tajikistan, Kyrgyzstan.
<i>SW Asia</i>	Afghanistan, western Pakistan (Baluchistan, Federally Administered Tribal Areas), Iran, Middle East, Arabian Peninsula, Turkey, Azerbaijan, Armenia, Georgia.
<i>Asia</i>	collective term for all above areas of Asia.
<i>Europe</i>	includes Ukraine, Belarus, Baltic republics.
<i>Africa</i>	includes Madagascar.
<i>N America</i>	includes C America south to Panama.
<i>S America</i>	south of Panama.
<i>Australasia</i>	Australia, New Zealand, Pacific Islands.
<i>Cosmopolitan</i>	collective term for a generally worldwide distribution.

Altitudes (elevation above sea level) are based on herbarium specimen data or records from credible observers. They are given to the nearest 100 m rounded up or down, with exceptional altitudes given in parentheses '()'. Likewise, flowering and fruiting times are based on specimens collected from Nepal, or on material from adjacent regions if these data are lacking and a note is provided to explain this. The short statement on the ecological preference of

each species and infraspecific taxon is mostly taken from herbarium specimen data. Currently these often lack detail, a reflection of the shortcomings of poor-quality data recorded by the past collectors of herbarium material, but these will improve with more field studies.

Supplementary information is given at the end of a taxon account discussing taxonomic issues, highlighting spot characters useful for identification, noting similar species that could cause confusion, and detailing the misapplication of names. Summary information is provided for ethnobotanical and other uses, but this is not intended to be exhaustive and is derived from secondary sources, such as *Plants and People of Nepal* and *A Compendium of Medicinal Plants of Nepal*.^{11, 12}

Abbreviations

Standard abbreviations for the International System of Units (SI) are used for measurements. Herbaria are cited using the standard abbreviation in *Index Herbariorum*.¹³ Other abbreviations used in the text include:

C	central.
ca. <i>circa</i>	about, approximately.
comb. nov. <i>combinatio nova</i>	new combination of name and epithet
dbh	diameter at breast height – measured on tree trunks at 1.3 m above the ground.
E	east, eastern.
et al. <i>et alia</i>	and others.
fig.	figure.
N	north, northern.
nom. cons. <i>nomen conservandum</i>	name officially conserved in ICBN. ¹⁴
nom. illegit. <i>nomen illegitimum</i>	illegitimate name, according to ICBN. ¹⁴
nom. inval. <i>nomen invalidum</i>	invalid name, according to ICBN. ¹⁴
nom. nud. <i>nomen nudum</i>	name lacking a description, or reference to an effectively published description, and so invalid according to ICBN. ¹⁴
nom. rej. <i>nomen rejiciendum</i>	name officially rejected in ICBN. ¹⁴
nom. superfl. <i>nomen superfluum</i>	name superfluous when published, and so illegitimate according to ICBN. ¹⁴
pl.	plate.
q.v. <i>quod vide</i>	which see.
S	south, southern.
s.l. <i>sensu lato</i>	for a taxon treated in a broad sense.
s.s. <i>sensu stricto</i>	for a taxon treated in a narrow sense.
sect.	section.
subfam.	subfamily.
subgen.	subgenus.
subsp.	subspecies.
subvar.	subvariety.
syn.	synonym
var.	variety.
W	west, western.
>	greater than
<	less than

References

1. Wu, Z.Y., Raven, P.H. & Hong, D.Y. (1994–ongoing). *Flora of China*, Science Press (Beijing) & Missouri Botanical Garden Press, St Louis [available online at flora.huh.harvard.edu/china].
2. Grierson, A.J.C., Long, D.G. & Noltie, H.J. (1983–2002). *Flora of Bhutan*, Royal Botanic Garden Edinburgh, Edinburgh.
3. The Angiosperm Phylogeny Group, M. W. Chase, M. J. M. Christenhusz, M. F. Fay, J. W. Byng, W. S. Judd, D. E. Soltis, D. J. Mabberley, A. N. Sennikov, P. S. Soltis, P. F. Stevens (2016). 'An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV'. *Bot. J. Linn. Soc.* 181: 1–20, doi.org/10.1111/boj.12385.
4. Hara, H., Stearn, W.T., Williams, W.T. & Chater, A.O. (1978, 1979, 1982). *An Enumeration of the Flowering Plants of Nepal*, 3 volumes, Trustees of the British Museum (Natural History), London.

5. Press, R., Shrestha, K.K. & Sutton, D.A. (2000). *Annotated Checklist of Flowering Plants of Nepal*, Natural History Museum: London & Tribhuvan University, Kathmandu [updated version available online at efloras.org].
6. Singh, A.P., Bista, M.S., Adhikari, M.K. & Rajbhandari, K.R. (2001). *Flowering Plants of Nepal (Phanerogams)*, HM Government of Nepal, Ministry of Forests, Department of Medicinal Plants, Kathmandu.
7. Brummit, R.K. & Powell, C.E. (1992). *Authors of Plant Names*, Royal Botanic Gardens, Kew, London [available online with revisions at www.ipni.org].
8. Bridson, G.D.R. & Smith, E.R. (1991). *Botanico-Periodicum-Huntianum*, ed. 2, Hunt Institute for Botanical Documentation, Pittsburgh.
9. Stafleu, F.A., Cowan, R.S. & Mennega, E. (1973–1988). *Taxonomic Literature*, ed. 2 (TL-2), Bonn, Scheltma & Holkema, Utrecht/Antwerpen; dr. W. Junk b.v., The Hague/Boston [available online at tl2.idcpublishers.info].
10. Harris, J.G. & Harris, M.W. (2001). *Plant Identification Terminology*, ed. 2, Spring Lake Publishing, Utah.
11. Manandhar, N.P. (2002). *Plants and People of Nepal*, Timber Press, Oregon.
12. Baral, S.R. & Kurmi, P.P. (2006). *A Compendium of Medicinal Plants in Nepal*, Mass Printing Press, Kathmandu.
13. Holmgren, P.K., Holmgren, N.H. & Barnett, L.C. (eds) (1990). *Index Herbariorum. Part 1: The Herbaria of the World*. ed. 8. New York Botanic Garden: New York. [available online with revisions at sweetgum.nybg.org/ih].
14. McNeill, J., Barrie, F.R., Burdet, H.M., Demoulin, V., Hawksworth, D.L., Marhold, K., Nicolson, D.H., Prado, J., Silva, P.C., Skog, J.E., Wiersema, J.H. & Turland, N.J. (eds) (2006). *International Code of Botanical Nomenclature (Vienna Code)*, Regnum Vegetabile 146. Gantner, Ruggell.